




Contents lists available at ScienceDirect

Review of Palaeobotany and Palynology

journal homepage: www.elsevier.com/locate/revpalbo

Multiproxy analysis on Indian wild ass (*Equus hemionus khur*) dung from Little Rann of Western India and its implications for the palaeoecology and archaeology of arid regions

Anil K. Pokharia^{a,*}, Sadhan K. Basumatary^{a,*}, Biswajeet Thakur^a, Swati Tripathi^a, H. Gregory McDonald^b, Deepika Tripathi^a, Pooja Tiwari^a, Eline Van Asperen^c, Michael Spate^d, Gaurav Chauhan^e, Mahesh G. Thakkar^e, Alka Srivastava^f, Shailesh Agarwal^a

^a Birbal Sahni Institute of Palaeosciences, 53 University Road, Lucknow 226007, Uttar Pradesh, India

^b Bureau of Land Management, Colorado State Office, 2850 Youngfield Street, Lakewood, CO 80215, USA

^c School of History, Classics and Archaeology, Newcastle University, Newcastle NE1 7RU, UK

^d Department of Archaeology, The University of Sydney, City Road, Camperdown, NSW 2006, Australia

^e Department of Earth and Environmental Science, KSKV Kachchh University, Bhuj 370001, India

^f Department of Botany, DG College, Kanpur 208001, India

ARTICLE INFO

Article history:

Received 14 December 2021

Received in revised form 18 May 2022

Accepted 27 May 2022

Available online 01 June 2022

Keywords:

Indian wild ass

Pollen

Dung midden

Arid region

Archaeology

Western India

ABSTRACT

The dung of the Indian wild ass was analyzed using biotic and abiotic proxies to determine its dietary habits in relation to the plant diversity and ecology in the arid region of western India. The presence of both micro and macrobotanical remains of Poaceae, Chenopodiaceae, and Fabaceae indicates they are the primary food plants of the wild ass. The continuous recovery of arboreal pollen taxa chiefly, *Prosopis*, *Acacia*, and *Ephedra* is indicative of dry thorny forest under semi-arid to arid conditions which display the existing vegetation and climate in the region. The recovery of marshy pollen taxa like Cyperaceae and Onagraceae along with *Arcella* indicates utilization of water-logged environments in the habitat. Spores of coprophilous fungi, *Sporormiella*, *Sordaria*, and *Podospora* are also present in the dung samples. The low value of stellate trichomes in winter dung samples reflects the seasonal migration of wild ass. Average $\delta^{13}\text{C}$ values ranging between -15.8% and -26.3% are indicative of a mixed diet of both C_3 and C_4 plants. The generated multiproxy data from dung samples can provide a reliable counterpart to modern data for the interpretation of the palaeoecology in relation to the palaeoherbivory and palaeodietary analysis in the region. This study also provides a basis to distinguish between wild and domesticated herbivores by analyzing coprolites and cultural sediments in archaeological sites.

© 2022 Published by Elsevier B.V.

1. Introduction

The Indian wild ass (*Equus hemionus khur*), an endangered subspecies of the Asian wild ass (Feh et al., 2002; Shah and Quershi, 2007; Kaczensky et al., 2016; Bennett et al., 2017) is today restricted to small areas in western India. Recent studies of the dung of endangered megaherbivores in tropical, subtropical and temperate regions of India to analyze their dietary patterns relative to the existing vegetation have been conducted as part of the modern analogue, species

conservation, and restoration strategy (Basumatary and McDonald, 2017; Tripathi et al., 2019; Basumatary et al., 2019, 2020, 2021). Building on these previous studies of modern herbivore dung as well as on coprolites of extinct species (Bryant Jr. and Larson, 1968; Bryant Jr., 1974; Riskind, 1970; Ghosh et al., 2003; Wood et al., 2008, 2021; Gill et al., 2009, 2013; Velazquez and Burry, 2012; Marinova et al., 2013; Gravendeel et al., 2014; Rawlence et al., 2016; Birks et al., 2018; Harrault et al., 2019; Van Geel et al., 2019) we have examined the dung of the Indian wild ass to better understand the species dietary preferences and ecology.

Likewise, the examination of botanical remains preserved in modern herbivore dung can also be applied to dung preserved in archaeological sites and can be a source of data for the interpretation of the palaeoecology, palaeovegetation, and climate in relation to the development of

* Correspondence authors at: Birbal Sahni Institute of Palaeosciences, 53 University Road, Lucknow 226007, Uttar Pradesh, India.

E-mail addresses: pokharia.anil@gmail.com (A.K. Pokharia), sbasumatary2005@yahoo.co.in (S.K. Basumatary).

human cultures and their impact on the local environment and fauna (Akeret et al., 1999; Baeten et al., 2018; Smith and Proctor, 2019; Rutunno et al., 2019; Dunseth et al., 2019; Fuks and Dunseth, 2021).

Previous palynological work on soils in northwestern India provides a basis to understand the modern pollen rain of the region and aid in reconstructing the palaeovegetation and climate (Singh et al., 1973). However, to date, no palynological work has been attempted on surface soil of Kachchh and its neighboring areas for developing a similar modern pollen analogue for palaeoecological interpretation of this region. The systematic study of modern pollen rain from surface samples of arid and semi-arid region is difficult due to the scarcity of depositional environments like forests, swamps, lakes, and wetlands that favor pollen preservation (Horowitz, 1992; Scott et al., 2022). Dung can serve as a complementary data source and counterpart to more traditional sources of pollen in the arid regions (Scott and Bousman, 1990; Scott and Cooremans, 1992; Scott and Cooremans, 1992; Scott and Vogel, 2000; Carrión, 2002; Pearson and Betancourt, 2002; Scott et al., 2004; Carrión et al., 2006; Marinova et al., 2013; Basumatary and McDonald, 2017; Basumatary et al., 2019). Dung as a substrate that preserves pollen in arid regions can also be sampled at different times of the year and thus can provide a basis to determine seasonal changes in plant diversity and phenology especially flowering and fruiting period of different plant species in this arid region. Such studies can augment scattered palaeoclimatic and palaeoecological work that has been carried out in the arid-semiarid environment of Gujarat (Prasad et al., 2007; Laskar et al., 2010; Pokharia et al., 2011, 2017, 2020; Farooqui et al., 2013; Prasad et al., 2014) in relation to the Indus and Harappan civilization (4600–3700 yr BP).

Stable carbon isotope analysis provides another approach to determine the preferences for C₃ and C₄ plants in an herbivore's diet in relation to the vegetation composition and climate in the region (Agrawal et al., 2012; Basu et al., 2015). Palynofacies analysis also aids in understanding their seasonal food habits. The palynofacies analysis of dung samples is subject to the dietary habits, the organic matter that the wild ass consumes (Hoen, 2000; Medeanic and Silva, 2010). It is observed that the seasonal vegetation is imparted as one of the feeding diets for the wild ass and hence the accumulation of dung primarily comprises the organic matter or vegetal remains that are eaten during the different seasons. Thus, the palynofacies in this way helps to ascertain the seasonal organic matter in dung sample (Hoen, 2000). In Mongolia the diet of the Asian wild ass is highly seasonal, and the species is primarily a grazer in the summer but a mixed feeder in winter (Sturm et al., 2017). Data on seasonal changes in elemental composition provides an indication of potential seasonal dietary stress resulting from changes in the availability of certain elements during part of the year (Levick and Rogers, 2008; Waldram et al., 2008; van der Waal et al., 2008, 2011; Ripple et al., 2015). Previously, no research has been conducted on the organic and inorganic contents of the dung of the Indian wild ass in relation to the vegetation composition and climate of western India. The main aim of this study is to conduct a multiproxy analysis of Indian wild ass dung in order to better understand the ecology of the species and provide a baseline to determine the palaeoherbivory, palaeoecology, and archaeology of this arid region.

1.1. Geographical distribution of hemionus

Grubb (1993) recognized three species of hemionus, *Equus hemionus*, *Equus kiang*, and *E. onager*. Only two species of hemionus, *E. hemionus* and *E. kiang* (Tibetan kiangs) were considered to represent the so-called Asiatic wild asses by Jónsson et al. (2014) and the onager was included as a subspecies of *E. hemionus*. Historically hemionus had a wide geographic range that included Mongolia, Central Asia, Anatolia, and Russia, but their current range is restricted to the border of Mongolia and China.

While the range of the *E. hemionus* was greater in the past based on remains recovered from Middle and Late Pleistocene deposits of the Purna Basin in Maharashtra (Thakuri et al., 2018), today its range is restricted

to the protected area in Little Rann of Gujarat, India as the population in Baluchistan became extinct over the past 40 years (Daniel, 1991). The spatial distribution of the Indian wild ass on the landscape is largely influenced by the distribution of food patches and water in this ecosystem.

1.2. Study area, vegetation, and wildlife

The present study was conducted in the Little Rann of Kachchh (LRK) and its surrounding in the Gujarat State of India (Fig. 1). The Little Rann is characterized by vast desiccated bare mudflats consisting of dark silt encrusted with salts. The LRK is dotted with around 74 elevated plateaus or islands. Due to the distinctiveness of the area and the existence of the endemic and endangered Indian wild ass the LRK was established as “Wild Ass Wildlife Sanctuary” in 1973 (Pasha et al., 2015). It is the largest wildlife sanctuary in India. The Little Rann of Kachchh is a seasonally wet-saline biome, with plants including grasses growing in a saline environment. The vegetation of the study areas includes dry thorny scrub forest composed of *Prosopis juliflora*, *Carissa spinosa*, *Ziziphus nummularia*, *Salvadora oleoides*, *Salvadora persica*, *Grewia*, *Lantana camera*, and *Ephedra*. The main ground cover vegetation in the region includes many species of Poaceae, Cyperaceae, Onagraceae, Chenopodiaceae, Malvaceae, and Amaranthaceae (Ishnava et al., 2011). The flora includes 372 species of flowering plants (Pandey, 2008).

Among wild herbivores, *E. hemionus khur* is commonly seen in the region (Fig. 2a). Other herbivores including, *Boselaphus tragocamelus* (Blue buck), *Gazella bennettii* (Indian gazelle), and *Rusa unicolor* (Sambar) are also commonly found in association with the Indian wild ass. The fauna also includes 307 species of birds (Pandey, 2008).

1.3. Climate and soil

The climate of the region is influenced by south-west monsoons. Annual rainfall is 125–400 mm with most of it occurring from July to September (Sinha and Goyal, 2006). During summer it is hot with a maximum temperature reaching 48 °C, while it is cold and dry in winter with the minimum temperature as low as 1 °C. The soil types of the study are primarily blackish sand with low amounts of organic matter. Agricultural fields and wasteland extend up to the borders of the LRK.

2. Material and methods

2.1. Field survey and sampling

During the winter (December–January) season of 2016–2017, AKP (first author) and team collected fifteen modern (fresh, 5–10 days old) dung samples (KW1–KW15). Dung selection was based on size and shapes so were considered to be from only adult population of wild ass (Fig. 2b). Similarly, in 2018–2019, GC (co-author) and team collected fifteen modern dung samples of the wild ass (KS1–KS15) during the summer (May–July) based on their size and shape. Each sample, consisting of approximately 200 g, was collected from in and around the Little Rann of Kachchh (Lat. 23°22'58.05"N – 23°35'29"N and Long. 70°52'26"E – 71°5'26.99"E). The outer surface of the collected dung samples was removed and the inner portion of the dung was packed separately in polythene bags to prevent contamination by pollen dispersed by wind and other possible contaminants before laboratory processing. Finally, the collected samples were mixed finely and taken to process for the different proxy analysis as per the sample's requirement.

2.2. Laboratory work

2.2.1. Macrobotanical analysis

For the macrobotanical analysis, 50 g of each of the 30 (15 per season) dung samples were gently boiled in 200 ml 5% KOH solution. After boiling, material was sieved through a 150 µm mesh prior to




Fig. 1. Location map showing the study area.

selection for study. The material was washed 2 to 4 times with distilled water and observed under stereo microscope (Leica Z6APO) and photographed with a Leica DFC295 camera (Fig. 3). The total of macroremains were counted and relative proportion (percentage) of each taxa was calculated. Identifications were made using published literature (Martin and Barkely, 1961; Musil, 1963) and reference specimens in the herbarium of the Birbal Sahni Institute of Palaeosciences (BSIP), Lucknow, India.

2.2.2. Microbotanical analysis

2.2.2.1. Pollen and non-pollen palynomorphs.

Thirty dung samples were processed for pollen and non-pollen palynomorphs (NPPs) using the standard acetolysis method (Erdtman, 1953). Samples were deflocculated using 10% solution of aqueous potassium hydroxide (KOH), treated with 40% hydrofluoric acid (HF) to dissolve silica, and acetolysis (9:1 anhydrous acetic anhydrite to concentrated sulfuric acid) for the removal of cellulose. The samples were washed twice with glacial acetic acid (GAA) and washed 3 to 4 times with distilled water. The samples were transferred to a 50% glycerol solution with a few drops of phenol to protect against microbial decomposition. 151 to 178 pollen and non-pollen palynomorphs (fungal spores, *Thecamoeba*, and algae) were counted from each sample to produce the palynomorph spectra. Results were plotted using TILIA software. The recovered pollen taxa were categorized as arboreal taxa (trees and shrubs), non-arboreal taxa (marshy and terrestrial herbs), fungal spores, *Thecamoeba*, and *Botryococcus*. For the identification of pollen grains, we consulted the reference slides at the Birbal Sahni Institute of Palaeosciences herbarium as well as published literature (Gupta and Sharma, 1986; Nayar, 1990; Farooqui et al., 2013) (Fig. 4).

2.2.3. Phytolith analysis

For phytolith analysis 30 samples from the summer and winter season were analyzed using the standard technique (Piperno, 2006). A 10 g sample of dried dung was treated with hydrochloric acid (HCl) to remove carbonates followed by removal of organic content by hydrogen peroxide (H_2O_2). Extraction of phytoliths was performed using a mixture of CdI_2 and KI (specific gravity 2.3) and placed on the slides using Canada Balsam. A total of 300–350 phytoliths were counted from each sample. Observation and microphotographs were done using an Olympus BX-61 microscope with DP-25 digital camera under $40\times$ magnification (Supplementary Fig. 1). Phytolith morphotypes were grouped according to the standard classification (Madella et al., 2005; Twiss et al., 1969; Mulholland and Rapp, 1992).

2.2.4. Palynofacies analysis

For palynofacies analysis, ~5 g of each sample was treated with 10% HCl to remove the carbonates from the dung samples. Repeated washing 3–4 times using distilled water was done to neutralize the sample and subsequently the residue was treated with 40% HF solution to dissolve the silicates. The samples were again thoroughly washed, sieved ($20\ \mu m$) and then the slides were prepared using polyvinyl alcohol and mounted using Canada Balsam (Tyson, 1995). No oxidizing agent was used during the processing of palynofacies analysis. The slides were then scanned under Olympus BH-2 microscope and photographs taken with DP-25 camera (Supplementary Fig. 1). The maceration process and literature consulted for the palynofacies analysis follows Traverse, 1994; Pocknall and Beggs, 1990; Batten, 1996; Roncaglia, 2004, and Sebag et al., 2006.




Fig. 2. Field photographs a. Close up of *Equus hemionus khur* in its natural habitat, b. A midden dung of *E. hemionus khur*, c. Practices of crop cultivation by the local people in and around the Little Rann of Katchch, d. A view of domesticated animals during drinking water time in and around the study areas.

2.2.5. PCA analysis

To obtain the vegetation and dietary differentiation in the dung samples procured from the Little Rann, Principal Component Analysis (PCA) was applied to the quantified pollen and phytolith frequencies analysis using CANOCO v.5 (Fig. 5) and the significance of the data was evaluated at p -value $\leq .05$ (Smilauer and Leps, 2014).

2.2.6. FESEM-EDS analysis

A Field Emission Scanning Electron Microscope (FESEM) with Energy Dispersive Spectroscopy (EDS) using FESEM (JEOL, JSM-7610F) equipped with EDS (EDAX, USA instrument) operated at 25 keV was used to determine the elemental composition of both summer and winter samples of the Indian wild ass dung (Fig. 6; Supplementary Table 1 and 2).

2.2.7. Stable carbon isotope analysis

For carbon isotope analysis ($\delta^{13}\text{C}_{\text{org}}$), ~ 1 g of the dung sample was decarbonized using HCl (5%) for 2 h in a water bath at 50 °C and the procedure was repeated three times. Subsequently the samples were centrifuged, rinsed repeatedly with deionized water until neutral pH was achieved and dried (Agrawal et al., 2012; Basu et al., 2015; Dubey et al., 2018; Ali et al., 2018). The de-carbonated samples were introduced into an Elemental Analyzer (Flash EA 2000 HT) through an auto sampler. Through the combustion CO_2 gas was produced and introduced into a Continuous Flow Isotope Ratio Mass Spectrometer (CFIRMS, MAT 253) coupled with Con-Flow IV interface for isotopic analysis. IAEA CH3 was used to calibrate the reference gas and carbon isotopic data has been reported against VPDB. International standards (IAEA CH3 and CH6) as well as internal standards (Sulfanilamide) were run to check the

accuracy for the CO_2 measurements with an external precision of $\pm 0.1\%$. Total organic carbon (TOC) was calculated from the peak area obtained from the sum of the integrated m/z 44, 45, and 46 signal measured in the CFIRMS (Fig. 7).

3. Results

3.1. Macrobotanical assemblage

Macrobotanical remains were recovered in samples from both seasons. Undigested seeds and leaves of Poaceae were predominant, with values upto 70% during summer and 50% during winter. The seeds of Chenopodiaceae and Amaranthaceae are also consistently present up to 20% during summer and 30% during winter in the macrobotanical assemblage. Arboreal taxa, especially the invasive species, *P. juliflora* along with *Acacia* pods, leaves, and seeds were also consistently present with values of 10% during summer and increase to 20% during the winter season.

3.2. Pollen and non-pollen palynomorph spectra (Fig. 8)

Poaceae–Cyperaceae–Prosopis–Grewia–Sporormiella–Sordaria–Arcella–Botryococcus assemblage: The dung samples (KS1–KS15) collected during the summer were characterized by the predominance of non-arboreal taxa (50.0%) over arboreal taxa (15.0%). Among non-arboreals, Poaceae are recorded at an average value of 22.0% in the pollen assemblage. The other non-arboreal taxa such as Cyperaceae, Chenopodiaceae, Amaranthaceae, and Malvaceae are also consistently present with values of 0.5% to 11.3%. Among the non-pollen




Fig. 3. Macrobotanical remains recovered from the *Equus hemionus khur dung* sample. a. *Panicum* (Poaceae), b. *Dactyloctenium* (Poaceae), c. and d. *Eleusine* (Poaceae), e. Poaceae seed, f. *Chenopodium* seed, g. *Chenopodium* germinating seed, h. and i. Amaranthaceae seed, j. *Celosia* germinating seed, k. Asteraceae seed, l, m. and n. Legume seed, o. Dicot leaf.

palynomorphs, the spores of coprophilous fungi, *Sporormiella*, *Podospora*, and *Sordaria* were recorded at 22.0% followed by non-coprophilous fungal spores (10.0%), *Arcella* (1.8%) and *Botryococcus* (1.3%).

Poaceae–Cyperaceae–Chenopodiaceae–Prosopis–Lantana–Sporormiella–Saccobolus–Arcella assemblage: The winter dung samples (KW1–KW15) were also characterized by the dominance of non-arboreal taxa

(60.6%) over arboreal taxa (10.6%). Among non-arboreals, Poaceae are present with an average value of 21.0% in the palynoassemblages. Other non-arboreal taxa such as Cyperaceae, Chenopodiaceae, Amaranthaceae, and Malvaceae are also consistently present but lower values of 0.5% to 12.8%. Among the non-pollen palynomorphs, the spores of coprophilous fungi *Sporormiella*, *Saccobolus*, and *Podospora* were most common at 21.2% followed by non-coprophilous fungal




Fig. 4. Pollen and non-pollen palynomorphs assemblage recovered from the *Equus hemionus khur dung* samples. a. and b. *Acacia* sp., c. *Grewia* sp., d. *Ephedra* sp., e. *Amaranthaceae*, f. *Chenopodiaceae*, g. *Malvaceae*, h. *Apiaceae*, i. *Solanaceae*, j. *Poaceae*, k. *Cyperaceae*, l. *Onagraceae*, m. *Arcella* sp., n. *Botryococcus* sp., o. *Glomus*, p. *Podospora* sp., q. *Sporormiella* sp., r. *Saccobolus*, s. *Pleospora* sp., t. *Bipolaris* sp., u. *Alternaria* sp.




Fig. 5. The PCA analysis of the pollen assemblages of *Equus hemionus khur dung* between summer and winter samples.

spores (6.6%). Both *Thecamoeba* (0.5%) and *Botryococcus* (0.5%) were consistently present at trace values or absent in the palynofacies in some samples.

3.3. Phytoliths spectra

In summer samples, the Poaceae are dominant in the assemblage across all samples. Among the Poaceae morphotypes, the frequency of grass silica short cells (GSSCs) was 53–69% of the sample. The GSSCs saddle morphotype was present with values of 17–23% followed by bilobates (10–16%), rondels (8–12%), trapezoids (1–9%), crosses (5–8%), and long saddles (1–6%). Elongate cells, including dendritic types and psilate types, were present at 1–5% and 1–4% respectively. Various multicell panels were also consistently encountered ranging from 2.5–12%. Among the non-Poaceae/arboreal phytolith types, the cylindrical sulcate (tracheary) morphology predominates with values of 12–20%, followed by acicular hair cells (1–8%) and stellate hair cells (1–6%). Cyperaceae morphotypes were recorded at 3–5% (Supplementary Fig. 2).

Similarly, in winter dung samples, the phytoliths of Poaceae predominated at 43–55%. Among GSSCs morphotype, the saddle type was recorded from 10 to 22%, followed by bilobates (11–16%), crosses (4–8%), trapezoids (2–8%), rondels (3–6%), and long saddles (2–5%). The elongate dendritic and elongate psilate are also recorded at 3–9% and 2–7% respectively. The cylindrical sulcate in arboreal types range from 13 to 18% followed by stellate hair types (3–9%) and acicular types (2–7%). A Cyperaceae type was also recorded at values of 2–5% (Supplementary Fig. 2).

3.4. Palynofacies spectra

The dung samples studied from the summer season are composed of a high percentage of brown degraded organic matter (DOM) (28–35%) with structured organic matter (SOM) ranging between 25 and 29%, stellate trichomes (18–20%), grass oxidized tissue (8–10%) and fungal remains (4–6%) respectively. The AOM (3–5%) and pollen/spore

(2–3%) occurrence is minimal. It is noted that in the SOM, the tracheid and cuticle vary in equitable proportion (Supplementary Fig. 3).

Similarly, the winter dung samples show an enhancement in all the components seen in the summer sample. The SOM has the highest representation of 34–36% followed by grass oxidized tissue (25–30%), brown degraded OM (20–25%), stellate trichome (10–15%), fungal remains (6–10%), pollen/spore (4–6%), and AOM (3–5%). In winter samples, the tracheids are better represented than the cuticle (Supplementary Fig. 3).

3.5. FESEM-EDS analysis data

The data generated from the FESEM-EDS elemental analysis of the summer dung samples recorded that the oxygen (O_2) content/level is 51.65 (weight %), followed by C, 37.3 (weight %), N, 4.42 (weight %), Si, 3.16 (weight %), Ca, 0.84 (weight %), Al, 0.78 (weight %), and K, 0.47 (weight %) (Fig. 6a) (Supplementary Table 1).

Similarly, the data recorded from the FESEM-EDS elemental analysis of the samples observed that the oxygen (O_2) content/level is 50.92 (weight %), followed by CK, 38.5 (weight %), N, 4.51 (weight %), Si, 2.89 (weight %), Al, 0.75 (weight %), and K, 0.37 (weight %) (Fig. 6b) (Supplementary Table 2).



3.6. Stable carbon isotope data

The average $\delta^{13}C$ values of summer dung samples varies from -15.8% to -26.3% . The TOC values were also recorded from 31.4% to 40.1%, respectively. Similarly, the average $\delta^{13}C$ values of winter dung samples were recorded from -28.1% to -30.9% . The TOC values vary from 37.89% to 44.3% and the TN% recorded from 1.1 to 1.91 respectively (Fig. 7), (Supplementary Table 3).

4. Discussion

The macro and microbotanical remains in Indian wild ass dung showed that the Poaceae are the primary food plants based on the dominance of both micro (pollen, phytoliths, and palynofacies) and macrobotanical remains. However, the recovery of pollen and macrobotanical remains of pods, seeds, and leaves of other families, especially of the Fabaceae, Chenopodiaceae, and Amaranthaceae, is indicative of their importance as food plants of the Indian wild ass. The regular presence of dicot phytoliths was marked in both the summer and winter samples, further indicating the importance of these plants to provide micronutrients. The arboreal taxa produce about twenty times fewer phytoliths than monocots (Albert et al., 2003). The regular presence of the chloridoid morphotype in the phytolith assemblage is indicative of the xerophytic nature in and around the study area. The Chenopodiaceae pollen consistently observed in the dung samples is indicative of the aridity in and around the study areas. Many chenopods utilize C_4 photosynthesis and are important ecologically in saline areas and in cold arid deserts (Pyankov et al., 2000). The presence of Cyperaceae pollen and phytoliths are indicative that despite the arid environment, the presence of water-logged habitat in the study area. The Indian wild ass requires drinking water at least once a day and is dependent on the availability of water bodies (Stubbe et al., 2005; Kaczensky et al., 2010).

The presence of pollen clumping, especially in Poaceae, Chenopodiaceae, and Amaranthaceae, observed in the palynofacies are indicative of a local origin and incorporation into the dung through the direct ingestion of the flowers and inflorescences. The regular recovery of Solanaceae and Apiaceae pollen along with Panicoidae morphotype phytoliths in the assemblage is a strong indicator of the anthropogenic activity in and around the study areas. This observation also indicates that the Indian wild ass is moving up to several kilometers for their food and dispersing into the surrounding human dominated landscape (Fig. 2c). The overall pollen data combining the summer and winter samples is



LEGENDS

X axes: Energy (Kilo Electron Volt)
 Y axes: Intensities

Fig. 6. FESEM-EDS analysis micrographs in *Equus hemionus khur dung* samples collected from summer (a) and winter (b).

indicative of dry thorny forest with large open areas under arid-to semi-arid conditions which corresponds with the current vegetation and climate in the region.

The abundance of palynofacies especially, DOM and SOM in the palynofacies assemblage is indicative of the composition of the vegetation and compliments the pollen data. The regular presence of stellate trichomes which are found on leaf surfaces (Wagner et al., 2004) have been observed in the assemblages. In the winter samples, it is observed that the structured OM remains are dominant in the palynofacies

assemblages, which could be a supplementary source of high nutrition for the Indian wild ass under the dry condition in the region. There is a lowering of stellate trichomes during winter that may reflect that either the Indian wild ass has moved to some nearby areas where it eats plants with a low content of stellate trichomes or the onset of environmental conditions during winter results in a reduction of stellar trichomes. The amorphous OM remains are appreciably similar to the winter samples which may reflect the environmental situations of the local topography.




Fig. 7. Stable carbon isotope analysis micrographs in *Equus hemionus khur* dung samples.

Among the fungal spores, the coprophilous fungi, especially *Sporormiella*, *Saccobolus*, and *Podospora* are dominant and consistently represented in both season's samples. The regular presence of *Sporormiella* and *Podospora* in the assemblage may simply reflect the cosmopolitan distribution of these two taxa. The non-coprophilous fungi, *Helminthosporium* and *Bipolaris*, which are pathogens of grassland plants, may have been incorporated in the dung through the ingestion of infected grasses and other herbs. The spores of the mycorrhizal fungi, *Glomus* along with hyphae (Fig. 4. o) are also consistently present in the assemblages and -is indicative of the impact of soil erosion (Van Geel, 1978; Anderson et al., 2011) in the region, resulting in dust settling on the vegetation, which is then ingested by the Indian wild ass with its food.

Determination of the nutrient element content in Indian wild ass dung provides information on the vegetation succession and biogeochemical cycle in arid and semi-arid regions. Plant species depend on different nutrient elements and their concentrations in the soil sediments (Koerselman and Meuleman, 1996; Aerts and Chapin, 1999;

Kroon, 2011). A total of 15 nutrient elements, both macro- and micro (C, N, O, Al, Si, Na, K, Cl, Ca, P, Mg, I, Fe, Mn, and Zn), have been recorded from the studied dung samples (Supplementary Table 1). Both macro and microelements required in the Indian wild ass diet can be provided inorganically or organically. The average stable carbon isotope analysis value of both summer and winter samples of the Indian wild ass dung is -20.6 , which indicates the presence of both C_3 and C_4 plants in their diet (Fig. 5). The stable isotope data complements the micro and macrobotanical remains identified in the dung samples.

4.1. PCA analysis for pollen and non-pollen palynomorphs

The pollen and non-pollen palynomorphs (NPPs) retrieved from the dung samples of Indian wild ass from the Rann of Kachchh during summer and winter were subjected to CANOCO 5.0 and based on response data it was found that the pollen and NPPs are compositional and have a gradient of 0.7 SD units long, hence linear method was recommended, and Principal Component Analysis (PCA) was performed on the present dataset. The variability explained by PCA1 and PCA2 comprises of 80% while the total variability of the first four principal components is 87% (Fig. 5).

The present study shows that during winter the amount of pollen is reduced as indicated by the PCA1. It can be observed that Solanaceae and Amaranthaceae are the highest recorded pollen taxa with the steepest increase of the values for corresponding pollen. These pollen taxa are closely correlated with other pollen types like *Ziziphus*, Cyperaceae, and Chenopodiaceae and to a certain extent with spores of coprophilous and non-coprophilous fungi like *Sporormiella* and *Alternaria*, respectively.

For the summer record of pollen and non-pollen palynomorphs (NPPs) in PCA1 it can be observed that *Prosopis*, Onagraceae, *Acacia*, *Grewia*, *Lantana*, Malvaceae, Poaceae, and *Ephedra* show the steepest increase of the values for corresponding pollen. These pollen taxa are indicative of enhanced arid conditions and low moisture conditions during any season and reflect very limited aquatic conditions. Consequently they can be said that as food plants they are high in availability in the Rann of Kachchh and hence their pronounced occurrence in the diet of the Indian wild ass in the region. These pollen taxa along with their enhanced occurrence are also correlated in the high probability of their presence in the dung as can be observed with the low angles in the same axis.

In the PCA2 axis, the NPPs with steepest increase are represented by *Botryococcus*, *Glomus*, *Tetraploa*, and *Pleospora* while the pollen and NPPs with increasing arrow heads are Amaranthaceae, *Sordaria*, *Bipolaris*, *Arcella*, *Saccobolus*, and *Helminthosporium*. Similarly, based on their correlation, we find that, these pollen types and NPP's are closely related, and they share a similar type of environmental preference and reflect the Rann of Kachchh environmental settings.

4.2. Indian wild ass dung and its implication for archaeology

Archaeological research on the rise of human civilization in western India has primarily focused on plant domestication (Pokharia et al., 2011, 2017). This has been complemented by studies on the domestication of cattle and sheep based on archaeozoological evidence (Patel, 2009). These traditional approaches can be complemented by the study of modern and fossil dung in relation to the soil sediments of archaeological sites in western India. While the materials found within animal dung are relevant to archaeology (Fig. 2d) the study of this component of a site is still wanting in Indian archaeological research.

The characteristics of the dung of animal species forms the basis of identification of archaeological dung deposits (Schelvis, 1992) which may represent various human activities on site, such as use of dung as a building material, for fuel and other purposes (Shahack-Gross, 2011). Small equid feces were encountered frequently during excavations in the Chehrabad salt mine in Iran (Askari et al., 2018). Given the potential




Fig. 8. Pollen and fungal spore spectra of the studied *Equus hemionus khur* dung samples.

preservation of equid and other feces at other archaeological sites, this study of Indian wild ass dung can serve as a model for coprolite as well as the soil sediments of the archaeological sites. The presence of spores of coprophilous fungi such as *Sporormiella*, *Podospora*, and *Sordaria* and differences in their relative abundance can result in distinct assemblages in the soil sediments in conjunction with pollen and non-pollen palynomorphs. For example, with an increase in the number of domestic cattle on the landscape, there should be a concurrent increase in dung fungal spores incorporated into the soil (Davis and Shafer, 2006) although there are many complicating factors for proper identification as the spore size varies (swelling or shrinking) due to chemical treatment (Van Geel et al., 2003; Van Geel and Aptroot, 2006; Van Asperen et al., 2021).

The presence of hemiones in archaeological sites may be due to either remains of wild individuals that were hunted, or potentially may be domesticated animals (Noble, 1969; Askari et al., 2018). There is evidence that other subspecies of hemiones such as the Persian onager (*Equus hemionus onager*) were domesticated elsewhere, such as at Chehrabad, an ancient salt mine in northwestern Iran (Askari et al., 2018). There is currently no evidence that the Indian subspecies or khur was domesticated. However, further study is needed to understand the interactions between past human populations and animals in India, including events such as extinction or domestication. The remains of equids have been reported from the Indus and Harappan archaeological site at Kanmer (5–3.5 ka BP) in the Kachchh District of Gujarat, along with wild and domesticated animals (Goyal et al., 2013). Elsewhere in Gujarat, it has been hypothesized that seminomadic pastoralists managed wild plants and herded animals from ca. 7000 BP onwards, culminating in the domestication of several millets and pulses (García-Granero et al., 2016). These types of habitat overlap partially with the current range of the Indian wild ass in Gujarat, and our data may have the potential for helping to better understand the types of habitats used by these past human populations and their interactions with wild herbivores.

These interactions may help contextualize potential remains of wild ass recovered from archaeological sites such as Shikarpur in Gujarat (Thomas et al., 2015) and SohrDamb Nal, Pakistan (Benecke and Neef, 2005). The archaeological site of Loteswar, located on the margin of a salty waste depression east of the Little Rann of Kachchh have also recorded hemione remains along with other faunal remains, all of which were considered wild species that were hunted in the older Microlithic deposits but hemiones are absent in the later Chalcolithic deposits which primarily contained remains of domesticated species (Patel, 2009).

5. Conclusions

Our multiproxy approach has contributed to understanding the diet of Indian wild ass in the arid-region of Kachchh, Gujarat, India and demonstrates the potential for utilizing these data to aid in examining animal–plant relationships and interpreting the interaction between humans and animals in the archaeological record. Due to the scarcity of water bodies on the landscape that normally accumulate and preserve pollen it is very difficult to study the modern pollen–vegetation relationship in arid regions. Our study of the dung of the Indian wild ass demonstrates that its pollen data reflects the thorny dry forest under arid condition and could serve as a reliable baseline for the interpretation of palaeovegetation and climate analysis. The consistent presence of *P. juliflora* in the dung of the wild ass supports previous studies (Pasha et al., 2015) that the spread of seeds of this invasive species is facilitated by its inclusion in the Indian wild ass diet. The negative impact by the exotic mesquite on grass availability in the Wild Ass Sanctuary (Sinha et al., 2009) reduces the availability of grasses which as shown here are a primary food source of the Indian wild ass. Consequently, recognition of the negative interaction between these two food sources of the Indian wild ass provides critical information for the management of this endangered species inhabiting a desert ecosystem where resources

can fluctuate widely. The stable carbon isotope data will be useful for the characterization of the herbivores coprolites as grazer or browsers in relation to the sedimentary soil in natural and archaeological sites in regional and global level. The analysis of coprolites can also be applied to the identification and differentiation between domesticated and wild herbivores in relation to pastoralism during early human civilization by both their analysis as well as the detailed examination of cultural sediments in western India and its neighboring areas and contribute to a better understanding of the role of domesticated and wild herbivores in the economies of these civilizations.

Declaration of Competing Interest

None.

Acknowledgements

We are thankful to Dr. Vandana Prasad, Director, Birbal Sahni Institute of Palaeosciences, Lucknow for laboratory facilities and permission to carry out this research work. We are thankful to Dr. Hukam Singh of BSIP for his help and information time to time. Thanks are also extended to the students of Institute of Rajasthan Studies, Udaipur, Rajasthan for their help in collecting dung samples. We thank both the reviewers for providing their valuable suggestions. We extend our appreciation to Dr. Bas van Geel for his insightful comments.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.revpalbo.2022.104700>.

References

- Aerts, R., Chapin, F.S., 1999. The mineral nutrition of wild plants revisited: a re-evaluation of processes and patterns. *Adv. Ecol. Res.* 30, 1–67.
- Agrawal, S., Sanyal, P., Sarkar, A., Jaiswal, M.K., Dutta, K., 2012. Variability of Indian monsoonal rainfall over the past 100 ka and its implication for C₃–C₄ vegetational change. *Quat. Res.* 77 (1), 159–170.
- Akeret, O., Haas, J.N., Leuzinger, U., Jacomet, S., 1999. Plant macrofossils and pollen in goat/sheep faeces from the Neolithic lake-shore settlement Arbon Bleiche 3, Switzerland. *The Holocene* 9 (2), 175–182.
- Albert, R.M., Bar-Yosef, O., Meignen, L., Weiner, S., 2003. Quantitative phytolith study of hearths from the Natufian and Middle Palaeolithic levels of Hayonim Cave (Galilee, Israel). *J. Archaeol. Sci.* 30 (4), 461–480.
- Ali, S.N., Dubey, J., Ghosh, R., Quamar, M.F., Sharma, A., Morthekai, P., Dimri, A.P., et al., 2018. High frequency abrupt shifts in the Indian summer monsoon since Younger Dryas in the Himalaya. *Sci. Rep.* 8 (1), 9287.
- Anderson, R.S., Homola, R.L., Davis, R.B., Jacobson, G., 2011. Fossil remains of the mycorrhizal fungal *Glomus fasciculatum* complex in post glacial lake sediments from Maine. *Can. J. Bot.* 6 (11), 2325–2328.
- Askari, Z., Mas-Coma, A.S., Bouwman, N., Boenke, T., Stöllner, A., Aalif, M., Rezaiana, G., et al., 2018. *Fasciola hepatica* eggs in paleofaeces of the Persian onager *Equus hemionus onager*, a donkey from Chehrabad archaeological site, dating back to the Sassanid Empire (224–651 AD), in ancient Iran. *Infect. Gen. Evol.* 62, 233–243.
- Baeten, J., Mees, P., Marinova, E., Dapper, M.D., Vos, D.D., Huyge, D., Strydonck, M.V., et al., 2018. Late Pleistocene coprolite from Qurta (Egypt) and the potential of the interdisciplinary research involving micromorphology, plant macrofossils and biomarker analysis. *Rev. Palaeobot. Palynol.* 259, 93–111.
- Basu, S., Agrawal, S., Sanyal, P., Mahato, P., Kumar, S., Sarkar, A., 2015. Carbon isotopic ratios of modern C₃–C₄ plants from the Gangetic Plain, India and its implications to paleovegetational reconstruction. *Palaeogr. Palaeoclimatol. Palaeoecol.* 440, 22–32.
- Basumatary, S.K., McDonald, H.G., 2017. Coprophilous fungi from dung of the Greater One-Horned Rhino in Kaziranga National Park, India and its implication to paleoherbivory and paleoecology. *Quat. Res.* 88, 14–22.
- Basumatary, S.K., Singh, H., McDonald, H.G., Tripathi, Swati, Pokharia, A.K., 2019. Modern botanical analogue of endangered Yak (*Bos mutus*) dung from India: plausible linkage with living and extinct megaherbivores. *PLoS One* 14 (3), e0202723.
- Basumatary, S.K., Singh, H., van Asperen, E.N., Tripathi, Swati, McDonald, H.G., Pokharia, A.K., 2020. Coprophilous and non-coprophilous fungal spores of *Bos mutus* dung from the Indian Himalaya: Implications to temperate paleoherbivory, paleoecological analysis. *Rev. Palaeobot. Palynol.* 277, 104208.
- Basumatary, S.K., Gogoi, R., Tripathi, Swati, Ghosh, Ruby, Pokharia, A.K., McDonald, H.G., Sherpa, N., van Asperen, E.N., et al., 2021. Red Panda feces from Eastern Himalaya as a modern analogue for palaeodietary and palaeoecological analyses. *Sci. Rep.* 11, 18312.
- Batten, D.J., 1996. Palynofacies and Palaeoenvironmental Interpretation. In: Jansonius, J., McGregor, D.C. (Eds.), *Palynology: Principles and Applications*. vol. 3. American Association of Stratigraphic Palynologists Foundation, College Station, TX, pp. 1011–1064.
- Benecke, N., Neef, R., 2005. Faunal and plant remains from Sohr Damb/Nal: a prehistoric site (c. 3500–2000 BC) in central Balochistan (Pakistan). In: Franke-Vogt U, Weishaar J, editors. *South Asian Archaeology 2003*. Aachen: Linden Soft. p. 81–91.
- Bennett, E.A., Champlot, S., Peters, J., Arbuckle, B.S., Guimaraes, S., Pruvost, M., et al., 2017. Taming the late Quaternary phylogeography of the Eurasian wild ass through ancient and modern DNA. *PLoS One* 12 (4), e0174216.
- Birks, H.H., van Geel, B., Fisher, D.C., Grimm, E.C., Kuijper, W.J., van Arkel, J., van Reenen, G.B.A., 2018. Evidence for the diet and habitat of two late Pleistocene mastodons from the Midwest, USA. *Quat. Res.* 1–21.
- Bryant Jr., V.M., 1974. Prehistoric diet in southwest Texas: the coprolite evidence. *Am. Antiq.* 39, 407–420.
- Bryant Jr., V.M., Larson, D.L., 1968. Pollen analysis of the Devil's Mouth Site, Val Verde County, Texas. In: Sorrow, B. (Ed.), *The Devil's Mouth Site: The Third Season*. Austin, USA: Papers of the Texas Archaeological Salvage Project. vol. 14, pp. 57–70.
- Carrión, J.S., 2002. A taphonomic study of modern pollen assemblages from dung and surface sediments in arid environments of Spain. *Rev. Palaeobot. Palynol.* 120, 217–232.
- Carrión, J.S., Scott, L., Arribas, A., Fuentes, N., Gil-Romera, G., Montoya, E., 2006. Pleistocene landscapes in central Iberia inferred from pollen analysis of hyena coprolites. *J. Quat. Sci.* 22, 191–202.
- Daniel, J.C., 1991. Ungulate conservation in India-Problems and prospects. *Appl. Anim. Behav. Sci.* 29, 348–359.
- Davis, O.K., Shafer, D.S., 2006. *Sporormiella* fungal spores, a palynological means of detecting herbivore density. *Palaeogr. Palaeoclimatol. Palaeoecol.* 237, 40–50.
- Dubey, J., Ghosh, R., Agrawal, S., Quamar, M.F., Morthekai, P., Sharma, R.K., Anupam, S.A., et al., 2018. Characteristics of modern biotic data and their relationship to vegetation of the Alpine zone of Chopta valley, North Sikkim, India: implications for palaeovegetation. *The Holocene* 28 (3), 363–376.
- Dunseth, Z.C., Fuks, D., Langgut, D., Weiss, E., Melamed, Y., Butler, D., Yan, X., et al., 2019. Archaeobotanical proxies and archaeological interpretation: a comparative study of phytoliths, seeds and pollen in dung pellets and refuse deposits at Early Islamic Shivta, Negev, Israel. *Quat. Sci. Rev.* 211, 166–185.
- Erdtman, G., 1953. *An Introduction to Pollen Analysis*. Chronica Botanica Company, Waltham, Mass, USA.
- Farooqui, A., Gaur, A.S., Prasad, V., 2013. Climate, vegetation and ecology during Harappan period: excavations at Kanjetar and Kaj, mid-Saurashtra coast, Gujarat. *J. Archaeol. Sci.* 40, 2631–2647.
- Feh, C., Shah, N., Rowen, M., Reading, R., Goyal, S.P., 2002. Status and action plan for the Asiatic wild Ass (*Equus hemionus*). In: Moehlman, P. (Ed.), *Equids: Zebras, Asses and Horse*. IUCN, Gland, Switzerland, pp. 62–70.
- Fuks, D., Dunseth, Z.C., 2021. Dung in the dumps: what we can learn from multi-proxy studies of archaeological dung pellets. *Veg. Hist. Archaeobot.* 30, 137–153.
- García-Granero, J.J., Lancelotti, C., Madella, M., Ajithprasad, P., 2016. Millets and herders: the origins of plant cultivation in semiarid North Gujarat (India). *Curr. Anthropol.* 57, 149–173.
- Ghosh, P., Bhattacharya, S.K., Sahni, A., Kar, R.K., Mohabey, D.M., Ambwani, K., 2003. Dinosaur coprolites from the Late Cretaceous (Maastrichtian) Lameta Formation of India: isotopic and other markers suggesting a C₃ plant diet. *Cretaceous Res.* 24, 743–750.
- Gill, J.L., Williams, J.W., Jackson, S.T., Lininger, K.B., Robinson, G.S., 2009. Pleistocene megafaunal collapse, novel plant communities, and enhanced fire regimes in North America. *Science* 326 (5956), 1100–1103.
- Gill, J.L., McLauchlan, K.K., Skibbe, A.M., Goring, S., Zirbel, C.R., Williams, J.W., 2013. Linking abundances of the dung fungus *Sporormiella* to the density of bison: implications for assessing grazing by megaherbivores in palaeorecords. *J. Ecol.* 101, 1125–1136.
- Goyal, P., Pokharia, A.K., Kharakwal, J.S., Joglekar, P., Rawat, Y.S., Osada, T., 2013. Subsistence system, palaeoecology, and ¹⁴C chronology at Kanmer, a Harappan site in Gujarat India. *Radiocarbon* 55 (1), 141–150.
- Gravendeel, B., Protopopov, A., Bull, I., Duijm, E., Gill, F., Nieman, A., Rudaya, N., et al., 2014. Multiproxy study of the last meal of a mid-Holocene Oygogor horse, Sakha Republic, Russia. *The Holocene* 24 (10), 1288–1296.
- Grubb, P., 1993. Order Perissodactyla. In: Wilson, D.E.; Reeder, D.A. (eds): *Mammal Species of the World. A taxonomic and geographic reference*. 2nd ed., pp. 369–372. - Smithsonian Institution Press, Washington and London.
- Gupta, H.P., Sharma, C., 1986. *Pollen Flora of North-West Himalaya*. Indian Association of Palynostratigraphers, Lucknow, India.
- Harrault, L., Milek, K., Jarde, E., Jeanneau, L., Derrien, M., Anderson, D.G., 2019. Faecal biomarkers can distinguish specific mammalian species in modern and past environments. *PLoS One* 14 (2), e0211119.
- Hoan, A.W., 2000. *The Use of Palynofacies Analysis in Archaeopalynology*. PhD Thesis University of Edinburgh, UK.
- Horowitz, A., 1992. *Palynology of Arid Lands*. Elsevier, Amsterdam, New York.
- Ishnava, K.V., Ramarao, J., Mohan, S.S., Kothari, L.L., 2011. Ecologically important and life supporting plants of Little Rann of Kachchh, Gujarat. *J. Ecol. Nat. Environ.* 3 (2), 33–38.
- Jónsson, H.M., Schubert, A., Seguin-Orlando, A., Ginolhac, L., Petersen, M., Fumagalli, A., Albrechtsen, B., et al., 2014. Speciation with gene flow in equids despite extensive chromosomal plasticity. *Proc. Natl. Acad. Sci.* 111 (52), 18655–18660.
- Kaczynsky, P., Dresley, V., Vetter, D., Otgonbayar, H., Walzer, C., 2010. Water use of Asiatic wildasses in the Mongolian Gobi. *Erfor. Biol. Resso. Mong.* 11, 291–298.
- Kaczynsky, P., Lkhagvasuren, B., Pereladova, O., Hemami, M., Bouskila, A., 2016. *Equus hemionus* ssp. khur. IUCN Red List of Threatened Species.
- Koerselman, W., Meuleman, A.F.M., 1996. The vegetation N:P ratio: a new tool to detect the nature of nutrient limitation. *J. Appl. Ecol.* 33, 1441–1450.

- Kroon, H., 2011. Large herbivores may alter vegetation structure of semi-arid savannas through soil nutrient mediation. *Oecologia* 165, 1095–1107.
- Laskar, A.H., Sharma, N., Ramesh, R., Jani, R.A., Yadav, M.G., 2010. Paleoclimate and paleovegetation of Lower Narmada Basin, Gujarat, Western India, inferred from stable carbon and oxygen isotopes. *Quat. Int.* 227, 183–189.
- Levick, S., Rogers, K., 2008. Patch and species specific responses of savanna woody vegetation to browser exclusion. *Biol. Conserv.* 141, 489–498.
- Madella, M., Alexandre, A., Ball, T., 2005. International code for phytolith nomenclature 1.0. *Ann. Bot.* 96, 253–260.
- Marinova, E., Ryan, P., Van Neer, W., Friedman, R., 2013. Animal dung from arid environments and archaeobotanical methodologies for its analysis: an example from animal burials of the Predynastic elite cemetery HK6 at Hierakonpolis, Egypt. *J. Environ. Archaeol.* 18 (1), 58–71.
- Martin, A.C., Barkely, W.D., 1961. Seed Identification Manual. University of California Press.
- Medeanic, S., Silva, M.B., 2010. Indicative value of non-pollen palynomorphs (NPPs) and palynofacies for palaeoreconstructions: Holocene Peat, Brazil. *Int. J. Coal Geol.* 84 (3–4), 248–257.
- Mulholland, S.C., Rapp, G., 1992. A morphological classification of grass silica-bodies. In: Rapp, G., Mulholland, S.C. (Eds.), *Phytolith Systematics. Advances in Archaeological and Museum Science*. vol. 1. Springer, Boston, MA, pp. 65–89.
- Musil, A., 1963. Identification of Crop and Weed Seeds, Agriculture Handbook No. 219. US Department of Agriculture, Washington, DC.
- Nayar, T.S., 1990. Pollen Flora of Maharashtra State, India. Today and Tomorrow's Printers & Publishers, New Delhi, India.
- Noble, D., 1969. The Mesopotamian Onager as a draught animal. In: Ucko, P.J., Dimbleby, G.W. (Eds.), *The Domestication and Exploitation of Plants and Animals*. Aldine Publishing Company, Chicago.
- Pandey, R.P., 2008. Studies on Phytodiversity of Little Rann of Kachchh with special reference to Wild Ass Sanctuary. *Gujarat. Bull. Bot. Surv. India* 50 (1–4), 39–50.
- Pasha, S.V., Satish, K.V., Reddy, C.S., Jha, C.S., 2015. Massive invasion of mesquite (*Prosopis juliflora*). *Nat. Acad. Sci. Lett.* 38, 271–273.
- Patel, A.K., 2009. Occupational histories, settlements, and subsistence in Western India: what bones and genes can tell us about the origins and spread of pastoralism? *Anthropozoologica* 44 (1), 173–188.
- Pearson, S., Betancourt, J.L., 2002. Understanding arid environments using fossil rodent middens. *J. Arid Environ.* 50, 499–511.
- Piperno, D.R., 2006. *Phytoliths: A Comprehensive Guide for Archaeologists and Paleoecologists*. Rowman. Altamira.
- Pocknall, D.T., Beggs, J.M., 1990. Palynofacies as a tool for the interpretation of depositional environments in the Waikato and Taranaki Basins, New Zealand. Proceedings of the 1989 New Zealand Oil Exploration Conference, Wellington (NZ). Petroleum and Geothermal Unit, Ministry of Commerce, pp. 250–258.
- Pokharia, A.K., Kharakwal, J.S., Rawat, Y.S., Osada, T., Nautiyal, C.M., Srivastava, A., 2011. Archaeobotany and archaeology at Kanmer, a Harappan site in Kachchh, Gujarat: evidence for adaptation in response to climatic variability. *Curr. Sci.* 100 (12), 1833–1846.
- Pokharia, A.K., Agnihotri, R., Sharma, S., Bajpai, S., Nath, J., Kumaran, R.N., Negi, B.C., 2017. Altered cropping pattern and cultural continuation with declined prosperity and extreme arid event at ~4.2ka BP: evidence from an Indus archaeological site Khirsara, Gujarat, western India. *PLoS One* 0185684.
- Pokharia, A.K., Kharakwal, J.S., Sharma, S., Spate, M., Tripathi, D., Dimri, A.P., Liu, X., et al., 2020. Variable monsoons and human adaptations: Archaeological and palaeoenvironmental records during the last 1400 years in north-western India. *The Holocene* 30 (9), 1332–1344.
- Prasad, V., Phartiyal, B., Sharma, A., 2007. Evidence of enhanced winter precipitation and the prevalence of a cool and dry climate during the mid to late Holocene in mainland Gujarat, India. *The Holocene* 17, 889–896.
- Prasad, Vandana, Farooqui, Anjum, Sharma, A., Phartiyal, Binita, Chakraborty, S., Bhandari, S., Raj, R., et al., 2014. Mid-late Holocene monsoonal variations from mainland Gujarat, India: a multi-proxy study for evaluating climate culture relationship. *Palaeogr. Palaeoclimatol. Palaeoecol.* 397, 38–51.
- Pyankov, V.I., Gunin, P.D., Tsoog, S., Black, C.C., 2000. C₄ plants in the vegetation of Mongolia: their natural occurrence and geographical distribution in relation to climate. *Oecologia* 123, 15–31.
- Rawlence, N.J., Wood, J.R., Bocherens, H., Rogers, K.M., 2016. Dietary interpretations for extinct megafauna using coprolites, intestinal contents and stable isotopes: Complimentary or contradictory? *Quat. Sci. Rev.* 142, 173–178.
- Ripple, W.J., Newsome, T.M., Wolf, C., Dirzo, R., Everatt, K.T., Galetti, M., Hayward, M.W., et al., 2015. Collapse of the world's largest herbivores. *Sci. Adv.* 1, e1400103.
- Riskind, D.H., 1970. Pollen analysis of human coprolites from Parida Cave. In: Alexander, B. (Ed.), *Archaeological Excavations at Parida Cave, Val Verde County, Texas*. Austin, USA: Papers of the Archaeological Salvage Project. vol. 19, pp. 89–101.
- Roncaglia, L., 2004. Palynofacies analysis and organic-walled dinoflagellate cysts as indicators of palaeo-hydrographic changes: an example from Holocene sediments in Skalaifjord, Faroe Islands. *Mar. Micropaleontol.* 50, 21–42.
- Rutunno, R., Marcuriam, Florenzano, A., Zerboni, A., di Lernia, S., 2019. Coprolite from rock shelter: hunter-gatherers "herding" Barbary sheep in the early Holocene Sahara. *J. Afr. Archaeol.* 17, 76–94.
- Schelvis, J., 1992. The identification of archaeological dung deposits on the basis of remains of predatory mites (Acari; Gamasida). *J. Archaeol. Sci.* 19, 677–682.
- Scott, L., Bousman, C.B., 1990. Palynological analysis of hyrax middens from Southern Africa. *Palaeogr. Palaeoclimatol. Palaeoecol.* 76, 367–379.
- Scott, L., Cooremans, B., 1992. Pollen in recent *Procavia* (Hyrax), *Petromus* (Dassie Rat) and bird dung in South Africa. *J. Biogeogr.* 19 (2), 205–215.
- Scott, L., Vogel, J.C., 2000. Evidence for environmental conditions during the last 20000 years in Southern Africa from ¹³C in fossil hyrax dung. *Glob. Planet. Change* 26, 207–215.
- Scott, L., Marais, E., Brook, G.A., 2004. Fossil hyrax dung and evidence of Late Pleistocene and Holocene vegetation types in the Namib Desert. *J. Quat. Sci.* 19, 1–4.
- Scott, L., Sobol, M., Neumann, F.H., Gil Romera, G., Fernández-Jalvo, Y., Bousman, C.B., Horwitz, L.K., van Aardt, A.C., 2022. Late Quaternary palaeoenvironments in the central semi-arid region of South Africa from pollen in cave, pan, spring, stream and dung deposits. *Quat. Int.* 614, 84–97.
- Sebag, D., Copard, Y., Di-Giovanni, C., Durand, A., Laignel, B., Ogier, S., Lallier-Verges, E., 2006. Palynofacies as useful tool to study origins and transfers of particulate organic matter in recent terrestrial environments: synopsis and prospect. *Earth Sci. Rev.* 79, 241–259.
- Shah, N., Quershi, Q., 2007. Social organization and determinants of spatial distribution of Khur (*Equus hemionus khur*). *Erforschung der biologischen Ressourcen der Mongolei (Halle/Saale)* 10, 189–200.
- Shahack-Gross, R., 2011. Herbivorous livestock dung: formation, taphonomy, method for identification, and archaeological significance. *J. Archaeol. Sci.* 38, 205–218.
- Smlauer, P., Lepš, J., 2014. *Multivariate Analysis of Ecological Data using CANOCO*. , p. 5 <https://doi.org/10.1017/cbo9781139627061>.
- Singh, G., Chopra, S.K., Singh, A.B., 1973. Pollen-rain from the vegetation of northwest India. *New Phytol.* 72, 191–206.
- Sinha, B.C., Goyal, S.P., 2006. Fuelwood plantation of *Prosopis juliflora* and its impact on the habitat of Indian wild ass, *Equus hemionus khur* in Little Rann of Kachchh, Gujarat. *Ann. Forest.* 14 (2), 350–354.
- Sinha, B.C., Goyal, S.P., Krauseman, P.R., 2009. How the use of mesquite impacts grass availability, Wild Ass Sanctuary, India. *Desert Plants* 25 (2), 3–9.
- Smith, A., Proctor, Hart, T.C., Stein, G.J., 2019. The burning issue of dung in archaeobotanical samples: a case study integrating macro-botanical remains, spherulites, and phytoliths to assess sample origin and fuel use at Tell Zeidan, Syria. *Veg. Hist. Archaeobot.* 28, 229–246.
- Stubbe, A., Stubbe, M., Batsajchan, N., Samjaa, R., Dorzderem, S., 2005. First results of wild ass research in the South Gobi Aymag/Mongolia in 2003 and 2004. *Erfor. Biol. Resso. Mong.* 9, 107–120.
- Sturm, M.B., Ganbaatar, O., Voigtand, C.C., Kaczynsky, P., 2017. Sequential stable isotope analysis reveals differences in dietary history of three sympatric equid species in the Mongolian Gobi. *J. Appl. Ecol.* 54, 1110–1119.
- Thakuri, N.G., Badam, G.L., Varade, A.M., 2018. Additional Quaternary faunal remains from the Middle-Late Pleistocene deposits of Purna Alluvial Basin, Maharashtra, Central India. *Curr. Sci.* 114 (7), 1402–1406.
- Thomas, P.K., Joglekar, P.P., Deshpande, A., Mukherjee, Pawankar, S.J., 2015. Harappan subsistence patterns with special reference to Shikarpur, a Harappan Site in Gujarat. *Man Environ* XX (2), 33–41.
- Traverse, A., 1994. *Sedimentation of Organic Particles*. Cambridge University Press, p. 557.
- Tripathi, S., Basumatary, S.K., Singh, Y.R., McDonald, H.G., Tripathi, D., Singh, L.J., 2019. Multiproxy studies on dung of endangered Sangai (*Rucervus eldii eldii*) and Hog deer (*Axis porcinus*) from Manipur, India: implications for paleoherbivory and paleoecology. *Rev. Palaeobot. Palynol.* 263, 85–103.
- Twiss, P.C., Suess, E., Smith, R.M., 1969. Morphological classification of grass Phytoliths 1. *Soil Sci. Soc. Amer. J.* 33, 109–115.
- Tyson, R.V., 1995. *Sedimentary organic matter: Organic facies and palynofacies*. Chapman and Hall, London.
- Van Asperen, E.N., Perotti, A., Baker, A., 2021. Coprophilous fungal spores: NPPs for the study of past megaherbivores. In: Marret, F., O'Keefe, J., Osterloff, P., Pound, M., Shumilovskikh, L. (Eds.), *Applications of Non-Pollen Palynomorphs: From Palaeoenvironmental Reconstructions to Biostratigraphy*. The Geological Society, London, Special Publications vol. 511, pp. 245–267.
- Van der Waal, C., Kool, A., Meijer, S.S., Kohi, E., Heitkoing, I.M., de Boer, W.F., van Langevelde, F., et al., 2008. The ecological implications of a Yakutian mammoth's last meal. *Quat. Res.* 69, 361–376.
- Van der Waal, C., Kool, A., Meijer, S.S., Kohi, E., Heitkönig, I.M.A., de Boer, W.F., van Langevelde, F., et al., 2011. Large herbivores may alter vegetation structure of semi-arid savannas through soil nutrient mediation. *Oecologia* 165, 1095–1107.
- Van Geel, B., 1978. A palaeoecological study of Holocene peat bog sections in Germany and the Netherlands, based on the analysis of pollen, spores and macro- and microscopic remains of fungi, algae, cormophytes and animals. *Rev. Palaeobot. Palynol.* 25, 1–120.
- Van Geel, B., Aptroot, A., 2006. Fossil Ascomycetes in Quaternary deposits. *Nova Hedwigia* 82, 313–329.
- Van Geel, B., Buurman, J., Brinkkemper, O., Schelvis, J., Aptroot, A., Vanreenen, G., Hakbijl, T., 2003. Environmental reconstruction of a Roman period settlement site in Uitgeest (The Netherlands), with special reference to coprophilous fungi. *J. Archaeol. Sci.* 30, 873–883.
- Van Geel, B., Langevelde, B.W., Mol, D., van der Knaap, P.W.O., van Leeuwen, J.F.N., 2019. Pollen and spores from molar folds reflect food choice of late Pleistocene and Early Holocene herbivores in The Netherlands and the adjacent North Sea area. *Quat. Sci. Rev.* 225, 106030.
- Velazquez, N.J., Burry, L.S., 2012. Palynological analysis of Lama guanicoe modern feces and its importance for the study of coprolites from Patagonia, Argentina. *Rev. Palaeobot. Palynol.* 184, 14–23.
- Wagner, G.J., Wang, E., Shepherd, R.W., 2004. New approaches for studying and exploiting an old protuberance, the plant trichome. *Ann. Bot.* 93, 3–11.

- Waldram, M.S., Bond, W.J., Stock, W.D., 2008. Ecological engineering by a mega-grazer: White Rhino impacts on a South African savanna. *Ecosystems* 11, 101–112.
- Wood, J.R., Rawlence, N.J., Rogers, G.M., Austin, J.J., Worthy, T.H., Cooper, A., 2008. Coprolite deposits reveal the diet and ecology of the extinct New Zealand megaherbivore moa (Aves, Dinornithiformes). *Quat. Sci. Rev.* 27, 2593–2602.
- Wood, J.R., Vermeulen, M.J., Bolstridge, N., Briden, S., Cole, T.L., Rivera-Perez, J., Shepherd, L.D., et al., 2021. Mid-Holocene coprolite from southern New Zealand provide new insights into the diet and ecology of the extinct little bush moa (*Anomalopteryx didiformis*). *Quat. Sci. Rev.* 263, 106992.